

The instruction on use of medicinal product
(for patients)

LIV-52[®] tablets

Composition

Active ingredients: Each tablet contains powders: Capparis spinosa 65mg, Cichorium intybus 65mg, Mandur bhasma 33mg, Solanum nigrum 32mg, Terminalia arjuna 32mg, Cassia occidentalis 16mg, Achillea millefolium 16mg, Tamarix gallica 16mg.

Processed over steam of an extract from a mix of medicinal raw materials: Eclipta alba, Phyllanthus amarus, Boerhaavia diffusa, Tinospora cordifolia, Raphanus sativus, Emblica officinalis, Plumbago zeylanica, Embelia ribes, Terminalia chebula, Fumaria officinalis.

Other ingredients: magnesium stearate, microcrystalline cellulose, croscarmellose sodium, colloidal silicon 4-oxide.

Pharmacotherapeutic group

Herbal preparation. Hepatoprotective and bile-expelling preparation.

Pharmacological properties

LIV-52 protects and restores hepatic parenchyma, raises functional efficiency of liver, accelerates regeneration and provides faster normalization of functions of a liver during infectious hepatitis. By its antiperoxidative activity, LIV-52 prevents the destruction of a cellular membrane. Normalizes level of cytochrome P-450. Facilitates rapid elimination of acetaldehyde and ensures protection from alcohol-induced hepatic damage. Preparation diminishes the lipotropic activity and prevents fatty infiltration of the liver in chronic alcoholism. As a daily health supplement, preparation improves appetite, the digestion and assimilation processes, and promotes weight gain.

Indications

1. In the prevention and treatment of:
 - viral hepatitis;
 - alcoholic liver disease;
 - pre-cirrhotic conditions and early cirrhosis;
 - protein energy malnutrition;
 - loss of appetite;
 - radiation and chemotherapy-induced liver damage.
2. As an adjuvant with hepatotoxic drugs.
3. A valuable adjuvant during convalescence and prolonged illness.

Contraindications

Hypersensitivity to product components.

Drug interactions

At simultaneous use:

Reduces level of ibuprofen in blood;

Reduces appropriating of tetracycline and doxycycline.

Special warnings

None.

Directions for use and dosage

Tuberculosis treatment:

2 tablets orally 2 times a day (together with anti-tuberculosis drugs)

Acute viral hepatitis:

children: 1 tablets 3 times a day;

adults: 2 tablets 3 times a day.

Cirrhosis:

2 tablets 3 times a day during 6-12 months.

Chronic hepatitis:

2 tablets 3 times a day during 6 months.

Chronic alcoholism of a liver:

2 tablets 3 times a day during 1 month, then 1 tablet 3 times a day during 6-12 months.

Anorexia and loss of weight:

children: 1 tablets 3 times a day;

adults: 2 tablets 3 times a day.

As a daily prevention:

2 tablets 2 times a day.

Adverse effects

Not reported.

Presentation

100 tablets in hermetic packages. One hermetic package with the instruction on use is put in cardboard packing.

Storage conditions

Store below +25°C in a place away from direct sunlight. Keep out of the reach of children. Do not refrigerate.

Expiry date

Three years from the date of manufacture.

Not to use after expired date.

Release form drugstores

Under the recipe.



Manufactured by
«The Himalaya Drug Company»
Makali, Bangalore 562 123, India
www.himalayahealthcare.com
® - registered trade mark



Official distributor in Azerbaijan
«TETRADADA» LTD.
AZ1102; 14, 20 Yanvar street; Baku, Azerbaijan
Tel.: (+994 12) 431-59-24, 431-05-41
Fax: (+994 12) 430-80-51
E-mail: info@tetradada-az.com
www.tetradada-az.com